

DAFTAR PUSTAKA

- Abraham, I. and Supriyati, Y. (2022) 'DESAIN KUASI EKSPERIMEN DALAM PENDIDIKAN: LITERATUR REVIEW', *Jurnal Ilmiah Mandala Education*, 8(3), pp. 2476–2482.
- Alloubani, A., Saleh, A. and Abdelhafiz, I. (2018) 'Hypertension and diabetes mellitus as a predictive risk factors for stroke', *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 12(4), pp. 577–584. Available at: <https://doi.org/10.1016/j.dsx.2018.03.009>.
- Anggriani, A. et al. (2018) 'PENGARUH ROM (Range of Motion) TERHADAP KEKUATAN OTOT EKSTREMITAS PADA PASIEN STROKE NON HEMORAGIC', *Jurnal Riset Hesti Medan Akper Kesdam I/BB Medan*, 3(2), p. 64. Available at: <https://doi.org/10.34008/jurhesti.v3i2.46>.
- Aries, A.M. et al. (2021) 'Sensory Stimulation of the Foot and Ankle Early Post-stroke: A Pilot and Feasibility Study', *Frontiers in Neurology*, 12. Available at: <https://doi.org/10.3389/fneur.2021.675106>.
- Bernhardt, J. et al. (2017) 'Early rehabilitation after stroke', *Current Opinion in Neurology*, 30(1), pp. 48–54. Available at: <https://doi.org/10.1097/WCO.0000000000000404>.
- Borstad, A. et al. (2022a) 'Tactile Sensation Improves Following Motor Rehabilitation for Chronic Stroke: The VIGoROUS Randomized Controlled Trial', *Neurorehabilitation and Neural Repair*, 36(8), pp. 525–534. Available at: <https://doi.org/10.1177/15459683221107893>.
- Borstad, A. et al. (2022b) 'Tactile Sensation Improves Following Motor Rehabilitation for Chronic Stroke: The VIGoROUS Randomized Controlled Trial', *Neurorehabilitation and Neural Repair*, 36(8), pp. 525–534. Available at: <https://doi.org/10.1177/15459683221107893>.
- Borstad, A. et al. (2022c) 'Tactile Sensation Improves Following Motor Rehabilitation for Chronic Stroke: The VIGoROUS Randomized Controlled Trial', *Neurorehabilitation and Neural Repair*, 36(8), pp. 525–534. Available at: <https://doi.org/10.1177/15459683221107893>.
- Chang, J.C. (2020) 'Stroke Classification: Critical Role of Unusually Large von Willebrand Factor Multimers and Tissue Factor on Clinical Phenotypes Based on Novel "Two-Path Unifying Theory" of Hemostasis', *Clinical and Applied Thrombosis/Hemostasis*, 26, p. 107602962091363. Available at: <https://doi.org/10.1177/1076029620913634>.

- Chen, L. et al. (2025) 'Effects of tactile feedback in post-stroke hand rehabilitation on functional connectivity and cortical activation: an fNIRS study', *Biomedical Optics Express*, 16(2), p. 643. Available at: <https://doi.org/10.1364/BOE.541820>.
- Chen, X. et al. (2018) 'Neonatal Exposure to Low-Dose (1.2%) Sevoflurane Increases Rats' Hippocampal Neurogenesis and Synaptic Plasticity in Later Life', *Neurotoxicity Research*, 34(2), pp. 188–197. Available at: <https://doi.org/10.1007/s12640-018-9877-3>.
- Cuppone, A.V., Semprini, M. and Konczak, J. (2018) 'Consolidation of human somatosensory memory during motor learning', *Behavioural Brain Research*, 347, pp. 184–192. Available at: <https://doi.org/10.1016/j.bbr.2018.03.013>.
- Dubuque, E.M., Collins, L. and Dubuque, M.L. (2021) 'Improving Performance Covertly and Remotely with Tactile Stimulation', *Behavior Analysis in Practice*, 14(1), pp. 203–207. Available at: <https://doi.org/10.1007/s40617-020-00493-0>.
- Faridah, A.A. et al. (2022) 'The Effectiveness of Range of Motion (ROM) on Increasing Muscle Strength in Stroke Patients: Literature Review', *Nursing and Health Sciences Journal (NHSJ)*, 2(2), pp. 137–142. Available at: <https://doi.org/10.53713/nhs.v2i2.118>.
- Frizzell, T.O. et al. (2022) 'Imaging functional neuroplasticity in human white matter tracts', *Brain Structure and Function*, 227(1), pp. 381–392. Available at: <https://doi.org/10.1007/s00429-021-02407-4>.
- Fuentes, B. (2018) 'Antidiabéticos en prevención de ictus en pacientes con diabetes tipo 2. El punto de vista del neurólogo', *Medicina Clínica*, 150(7), pp. 275–281. Available at: <https://doi.org/10.1016/j.medcli.2017.09.012>.
- Govindarajan, P. et al. (2020) 'RETRACTED ARTICLE: Classification of stroke disease using machine learning algorithms', *Neural Computing and Applications*, 32(3), pp. 817–828. Available at: <https://doi.org/10.1007/s00521-019-04041-y>.
- Holzapfel, S.D. et al. (2019) 'Acute Effects of Assisted Cycling Therapy on Post-Stroke Motor Function: A Pilot Study', *Rehabilitation Research and Practice*, 2019, pp. 1–10. Available at: <https://doi.org/10.1155/2019/9028714>.
- Huynh, B.P. et al. (2023) 'Sensitivity to Change and Responsiveness of the Upper Extremity Fugl-Meyer Assessment in Individuals With Moderate to Severe Acute Stroke', *Neurorehabilitation and Neural Repair*, 37(8), pp. 545–553. Available at: <https://doi.org/10.1177/15459683231186985>.
- Johnson, D. et al. (2018) 'Interrater Reliability of Three Versions of the Chedoke Arm and Hand Activity Inventory', *Physiotherapy Canada*, 70(2), pp. 133–140. Available at: <https://doi.org/10.3138/ptc.2016-70>.

- Jurkowski, M.P. et al. (2020) 'Beyond the Hippocampus and the SVZ: Adult Neurogenesis Throughout the Brain', *Frontiers in Cellular Neuroscience*, 14. Available at: <https://doi.org/10.3389/fncel.2020.576444>.
- Karaca, O. and Kılınç, M. (2024) 'Sensory training combined with motor training improves trunk proprioception in stroke patients: a single-blinded randomized controlled trial', *Neurological Research*, 46(6), pp. 553–560. Available at: <https://doi.org/10.1080/01616412.2024.2337522>.
- Kim, Y.W. (2022) 'Update on Stroke Rehabilitation in Motor Impairment', *Brain & Neurorehabilitation*, 15(2). Available at: <https://doi.org/10.12786/bn.2022.15.e12>.
- Kisner, C., Colby, L.A. and Borstad, J. (2022) *Therapeutic Exercise Foundations and Techniques*. Eight. F A Davis Co.
- Kitai, K. et al. (2021) 'Evaluation of Intervention Effectiveness of Sensory Compensatory Training with Tactile Discrimination Feedback on Sensorimotor Dysfunction of the Hand after Stroke', *Brain Sciences*, 11(10), p. 1314. Available at: <https://doi.org/10.3390/brainsci11101314>.
- Kristensen, M.G.H., Busk, H. and Wienecke, T. (2022) 'Neuromuscular Electrical Stimulation Improves Activities of Daily Living Post Stroke: A Systematic Review and Meta-analysis', *Archives of Rehabilitation Research and Clinical Translation*, 4(1), p. 100167. Available at: <https://doi.org/10.1016/j.arrct.2021.100167>.
- Kwon, H., Kim, J. and Lee, M. (2020) 'Brain activation induced by different strengths of hand grasp: a functional magnetic resonance imaging study', *Neural Regeneration Research*, 15(5), p. 875. Available at: <https://doi.org/10.4103/1673-5374.268907>.
- Lee, K.E., Choi, M. and Jeoung, B. (2022) 'Effectiveness of Rehabilitation Exercise in Improving Physical Function of Stroke Patients: A Systematic Review', *International Journal of Environmental Research and Public Health*, 19(19), p. 12739. Available at: <https://doi.org/10.3390/ijerph191912739>.
- Lin, Y.-F. et al. (2021) 'Weakened Effective Connectivity Related to Electroacupuncture in Stroke Patients with Prolonged Flaccid Paralysis: An EEG Pilot Study', *Neural Plasticity*, 2021, pp. 1–10. Available at: <https://doi.org/10.1155/2021/6641506>.
- Marzola, P. et al. (2023) 'Exploring the Role of Neuroplasticity in Development, Aging, and Neurodegeneration', *Brain Sciences*, 13(12), p. 1610. Available at: <https://doi.org/10.3390/brainsci13121610>.
- Mihara, M. et al. (2021) 'Effect of Neurofeedback Facilitation on Poststroke Gait and Balance Recovery', *Neurology*, 96(21). Available at: <https://doi.org/10.1212/WNL.0000000000011989>.
- Murphy, S.JX. and Werring, D.J. (2020) 'Stroke: causes and clinical features', *Medicine*, 48(9), pp. 561–566. Available at: <https://doi.org/10.1016/j.mpmed.2020.06.002>.

- Nam, K.E. et al. (2019) 'When does spasticity in the upper limb develop after a first stroke? A nationwide observational study on 861 stroke patients', *Journal of Clinical Neuroscience*, 66, pp. 144–148. Available at: <https://doi.org/10.1016/j.jocn.2019.04.034>.
- Ostry, D.J. and Gribble, P.L. (2016) 'Sensory Plasticity in Human Motor Learning', *Trends in Neurosciences*, 39(2), pp. 114–123. Available at: <https://doi.org/10.1016/j.tins.2015.12.006>.
- Patil, S. et al. (2022) 'Detection, Diagnosis and Treatment of Acute Ischemic Stroke: Current and Future Perspectives', *Frontiers in Medical Technology*, 4. Available at: <https://doi.org/10.3389/fmedt.2022.748949>.
- Piepmeyer, A.T. and Etnier, J.L. (2015) 'Brain-derived neurotrophic factor (BDNF) as a potential mechanism of the effects of acute exercise on cognitive performance', *Journal of Sport and Health Science*, 4(1), pp. 14–23. Available at: <https://doi.org/10.1016/j.jshs.2014.11.001>.
- Pinar, C. et al. (2017) 'Revisiting the flip side: Long-term depression of synaptic efficacy in the hippocampus', *Neuroscience & Biobehavioral Reviews*, 80, pp. 394–413. Available at: <https://doi.org/10.1016/j.neubiorev.2017.06.001>.
- Pratiwi, R.M. (2023) 'The Effect of Cylindrical Grip ROM Exercise Therapy on Finger Grip Muscle Strength in The Upper Extremities of Post Stroke Patients', *Jurnal Ners dan Kebidanan (Journal of Ners and Midwifery)*, 10(3), pp. 329–338. Available at: <https://doi.org/10.26699/jnk.v10i3.ART.p329-338>.
- Pu, L. et al. (2023) 'Projected Global Trends in Ischemic Stroke Incidence, Deaths and Disability-Adjusted Life Years From 2020 to 2030', *Stroke*, 54(5), pp. 1330–1339. Available at: <https://doi.org/10.1161/STROKEAHA.122.040073>.
- Rahayu, K.I.N. (2015) 'PENGARUH PEMBERIAN LATIHAN RANGE OF MOTION (ROM) TERHADAP KEMAMPUAN MOTORIK PADA PASIEN POST STROKE DI RSUD GAMBIRAN The Influence of Range of Motion Exercise to Motor Capability of Post-Stroke Patient at the Gambiran Hospital', *Jurnal Keperawatan*, 6(2), pp. 102–107.
- Rani, D.M. et al. (2022) *Anatomi Fisiologi Tubuh Manusia*. Medan: Yayasan Kita Menulis.
- Rexrode, K.M. et al. (2022) 'The Impact of Sex and Gender on Stroke', *Circulation Research*, 130(4), pp. 512–528. Available at: <https://doi.org/10.1161/CIRCRESAHA.121.319915>.
- Sanz-Esteban, I. et al. (2021) 'Cortical activity during sensorial tactile stimulation in healthy adults through Vojta therapy. A randomized pilot controlled trial', *Journal of NeuroEngineering and Rehabilitation*, 18(1), pp. 1–13. Available at: <https://doi.org/10.1186/s12984-021-00824-4>.

- Schnakers, C., Magee, W.L. and Harris, B. (2016) 'Sensory stimulation and music therapy programs for treating disorders of consciousness', *Frontiers in Psychology*, 7(MAR), pp. 1–6. Available at: <https://doi.org/10.3389/fpsyg.2016.00297>.
- Serrada, I., Hordacre, B., and Hillier, S.L. (2019) 'Does Sensory Retraining Improve Sensation and Sensorimotor Function Following Stroke: A Systematic Review and Meta-Analysis', *Front. Neurosci.* 13:402. doi: 10.3389/fnins.2019.00402.
- Setyopranoto, I. et al. (2019) 'Prevalence of stroke and associated risk factors in sleman district of Yogyakarta Special Region, Indonesia', *Stroke Research and Treatment*, 2019. Available at: <https://doi.org/10.1155/2019/2642458>.
- Shahid, J., Kashif, A. and Shahid, M.K. (2023) 'A Comprehensive Review of Physical Therapy Interventions for Stroke Rehabilitation: Impairment-Based Approaches and Functional Goals', *Brain Sciences*, 13(5), p. 717. Available at: <https://doi.org/10.3390/brainsci13050717>.
- Stolwyk, R.J. et al. (2024) 'Post-stroke Cognition is Associated with Stroke Survivor Quality of Life and Caregiver Outcomes: A Systematic Review and Meta-analysis', *Neuropsychology Review* [Preprint]. Available at: <https://doi.org/10.1007/s11065-024-09635-5>.
- Sutopo, H., Murti, B. and Demartoto, A. (2023) 'Effect of Robot Assistive Therapy on Hand Functional Ability in Stroke Patients: Meta-Analysis', *Journal of Epidemiology and Public Health*, 8(2), pp. 213–222. Available at: <https://doi.org/10.26911/jepublichealth.2023.08.02.06>.
- Syah, L.O.M.G. et al. (2020) 'Biofeedback Exercise Lebih Baik Daripada Active Assisted Exercise Untuk Meningkatkan Kinerja Otot Bahu Pada Fungsional Meraih Posisi 900 Fleksi Bahu Pasien Pasca Stroke', *Sport and Fitness Journal*, 8(1), p. 29. Available at: <https://doi.org/10.24843/spj.2020.v08.i01.p05>.
- Syukriyah, S.R.Z.M.D. and Fauzi, A. Al (2024) 'Hypertension as a risk factor in stroke: An overview', *World Journal of Advanced Research and Reviews*, 21(1), pp. 2370–2372. Available at: <https://doi.org/10.30574/wjarr.2024.21.1.0204>.
- Wang, Y. et al. (2021) 'A robot-aided visuomotor wrist training induces gains in proprioceptive and movement accuracy in the contralateral wrist', *Scientific Reports*, 11(1), p. 5281. Available at: <https://doi.org/10.1038/s41598-021-84767-9>.
- West, K. et al. (2021) "'People Associate Us with Movement so It's an Awesome Opportunity": Perspectives from Physiotherapists on Promoting Physical Activity, Exercise and Sport', *International Journal of Environmental Research and Public Health*, 18(6), p. 2963. Available at: <https://doi.org/10.3390/ijerph18062963>.
- Van de Winckel, A. et al. (2020) 'Exploratory study of how Cognitive Multisensory Rehabilitation restores parietal operculum connectivity and improves upper limb

movements in chronic stroke', *Scientific Reports*, 10(1), p. 20278. Available at: <https://doi.org/10.1038/s41598-020-77272-y>.

Zandvliet, S.B. et al. (2020) 'Is Recovery of Somatosensory Impairment Conditional for Upper-Limb Motor Recovery Early After Stroke?', *Neurorehabilitation and Neural Repair*, 34(5), pp. 403–416. Available at: <https://doi.org/10.1177/1545968320907075>.